

Higher rates of concussion following COVID-19 infection in high school athletes

Garrett S Bullock ¹, Carolyn A Emery ², Vicki R Nelson,³ Albert Prats-Uribe,⁴ Robert Gill Gilliland,⁵ Charles A Thigpen,⁶ Ellen Shanley ^{5,6}

¹Department of Orthopaedic Surgery & Rehabilitation, Wake Forest University School of Medicine, Winston-Salem, North Carolina, USA

²Sport Injury Prevention Research Centre, University of Calgary, Calgary, Alberta, Canada

³Sports Medicine, Prisma Health, Greenville, South Carolina, USA

⁴Pharmaco Epi-Device Unit, University of Oxford, Oxford, UK

⁵Sports Medicine, ATI Physical Therapy, Bolingbrook, Illinois, USA

⁶Clinical Excellence, ATI Physical Therapy, Greenville, Illinois, USA

Correspondence to

Dr Garrett S Bullock, Department of Orthopaedic Surgery & Rehabilitation, Wake Forest University School of Medicine, Winston-Salem, North Carolina, USA; garrettbullock@gmail.com

Accepted 31 January 2023
Published Online First
8 February 2023

ABSTRACT

Objective To compare concussion rates (CRs) over one academic year in high school athletes with and without a COVID-19 infection prior to concussion.

Methods Illness and concussion were prospectively reported for male and female high school athletes across six states over one academic year in the Players Health Rehab surveillance system. Concussion was truncated to 60 days following recovery and return to sport from COVID-19. CRs were estimated per 1000 athletes per academic year and stratified by those who tested positive for COVID-19 infection (with COVID-19) and those who did not (no COVID-19). Poisson regression analyses estimated rate ratio (RR) of concussion controlling for state, gender and an offset of the log athlete participation (with COVID-19 and no COVID-19).

Results Of 72 522 athletes, 430 COVID-19 infections and 1273 concussions were reported. The CR was greater in athletes who reported COVID-19 (CR=74.4/1000 athletes/year, 95% CI 49.6 to 99.3) compared with those who did not (CR=17.2, 95% CI 16.3 to 18.2). Athletes with recent COVID-19 had a threefold higher rate of concussion (RR=3.1, 95% CI 2.0 to 4.7).

Conclusion Athletes returning from COVID-19 had higher CRs than those who did not experience COVID-19. This may be related to ongoing COVID-19 sequelae or deconditioning related to reduced training and competition load during the illness and when returning to sport. Further research is needed to understand the association of recent COVID-19 infection and concussion in order to inform preventive strategies.

INTRODUCTION

Concussions in high school athletes are a serious problem. Almost 20% of all adolescents in the USA report experiencing a concussion during high school,^{1,2} with over 700 000 concussions sustained by high school athletes annually.³ Concussion rates (CRs) range from 2 to 20 concussions per 10 000 athletic exposures, and from 10% to 13% of all injuries suffered by high school athletes are concussions.^{3,4} Adolescent athletes may be at higher risk of sustaining a concussion compared with adult athletes,^{5,6} with more severe symptoms and potential developmental implications including cognitive dysfunction.⁷

Higher CRs are reported for girl athletes compared with boy athletes in gender comparable sports.^{8,9} These gender discrepancies may be resultant of differences in physiology,^{10,11} with likelihood of reporting concussions,¹² protective gear,⁸ player contact and strategy,¹³ ball to head

WHAT IS ALREADY KNOWN ON THE TOPIC

⇒ Concussions can have serious consequences, particularly in adolescents. It is currently unclear how COVID-19 infections may impact subsequent concussion risk in sport.

WHAT THIS STUDY ADDS

⇒ Athletes with COVID-19 infection had greater than threefold rates of concussion within 60 days following recovery from infection compared with athletes without COVID-19 infection.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ Additional research is warranted to explore why athletes with a recent COVID-19 infection are at higher risk of concussion. Clinical judgement or concern may guide further evaluation or a more cautious return to sport in athletes recovering from COVID-19.

ratio¹³ or practices.^{14,15} There are also differences in CRs between sports. Sports with high levels of player contact have greater exposure to potential head trauma, including both collisions and field and court sports.^{4,16}

SARS-CoV-2 and the resultant disease, COVID-19 infection, have significantly impacted adolescent health.^{17,18} COVID-19 infections can result in respiratory, cardiovascular, neurological and musculoskeletal system issues, long after acute COVID-19 has resolved.¹⁹ COVID-19 can also impair neurophysiological processes,²⁰ potentially resulting in impaired cognition.²¹ While adults report increased COVID-19 symptoms, adolescents report a high prevalence of symptoms, ranging from 4% to 80%.²²

Concussions have serious consequences including memory loss and behaviour issues,^{15,23} with adolescents being more vulnerable to downstream sequela due to their developing brains.⁵⁻⁷ While COVID-19 infection rates have been established and long-term physiological and neurological consequences have been identified,¹⁹ it is currently unknown how COVID-19 infections may impact subsequent concussion risk following illness recovery. This information would improve clinical examination, decision making, and athlete and parent education regarding returning to sport. Therefore, the purpose of this study was to compare CRs over one academic year in high school athletes with and without a COVID-19 infection prior to concussion.



© Author(s) (or their employer(s)) 2023. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Bullock GS, Emery CA, Nelson VR, et al. *Br J Sports Med* 2023;**57**:590–594.

METHODS

Study design

A prospective cohort study was conducted over one academic year to compare CRs in athletes who reported a COVID-19 infection prior to concussion and those who did not. The Strengthening the Reporting of Observational Studies in Epidemiology for Sport Injury and Illness Surveillance was used for reporting.²⁴

Patient and public involvement

Physicians, athletic trainers, physical therapists, former athletes, high school coaches and school administrators from multiple geographical regions, urban and rural communities, and different socioeconomic backgrounds were interviewed and gave input into the collection methods and electronic medical record dashboard. Findings from this concussion study were presented to these stakeholders along with one-page reports in lay language.

Equity, diversity and inclusion

High schools from multiple geographical regions, urban and rural communities, and different socioeconomic backgrounds were recruited for this study. Participants could report gender beyond binary terms. The author team included early, middle and late career researchers with balance from people who identify as male and female.

Outcomes

An illness was defined as a new physical complaint or disorder experienced by an athlete who was not related to an injury. Illnesses included health problems relating to physical complaints or removal or loss of vital elements (ie, dehydration, hypothermia or fever).^{24–26} A COVID-19 infection was defined as a positive test administered by a healthcare professional. COVID-19 screening was performed daily by the high school assigned athletic trainers for each athlete, prior to clearance for sports participation. Screening included measurement of body temperature, along with symptom reporting. Any athletes who had elevated body temperatures and/or reported COVID-like symptoms were tested for COVID-19. Athletes were quarantined within 24 hours following documentation of COVID-19 infection, with athletes being asymptomatic or symptomatic at the time of quarantine. Chronic pre-existing illnesses (eg, asthma) were not included unless there was an acute exacerbation requiring medical attention. Illness severity was calculated for overall time loss, with further illness severity stratified by 1–6 (minor), 7–27 (moderate) and 28+ (severe) days.²⁴

Concussion was defined as a traumatic brain injury, integrating clinical, pathological and biomechanical features.²⁷ Concussion was induced by contact to the head, neck or body that resulted in forces sustained to head, which resulted in rapid or delayed onset of neurological symptoms.²⁷ Potential concussions were evaluated on field and within the on-site training room. All potential concussions were referred to a sport physician for follow-up. The time frame whereby a concussion was deemed as following a COVID-19 infection was truncated to 60 days following illness recovery. Illness recovery was defined as a full return to training and/or competition following symptom resolution.^{24 25}

All athletes presenting with any illness reported to their school athletic trainer to verify the complaint. The athletic trainer documented the illness in the Players Health Rehab system. The documentation included the illness date, athlete gender, sport of participation, problem type, the participation status of the athlete, athlete phase of recovery and days until return to sport.

Data collection

Athlete illness data were recorded by the high school athletic trainer, including the date and duration of time loss during a team-sponsored practice or game. Sport coverage was determined through the proportion of high school students who participated in sport, not on high school or school district funding. High school sport participation per season was determined through rosters by each athletic trainer. The high school athletic trainers collected athlete participation each day. Internal validity data checks were performed by the regional athletic supervisor and the regional and national quality control supervisors on a quarterly basis.

Participants

High school athletes from six states (Alabama, Delaware, Illinois, Maryland, Michigan and Pennsylvania) were included. High schools had to participate in all three sport seasons (eg, fall, winter and spring) of the 2020–2021 academic year for inclusion.

Exposure

An athlete exposure (AE) was defined as one athlete participating in one practice or competition where a player was at risk of sustaining an injury.²⁸ AE was not calculated for days of school suspension, medical visits that were not musculoskeletal related (eg, dentist) or extramural school activities not related to sport. Exposure following illness was not recorded until the athlete had recovered from illness, which was defined as when an athlete fully returned to training and/or competition.^{24 25}

Explanatory variables

Self-identified gender, state and sport were identified as important covariables. Due to the large number of sports played, four categories were identified: collision (American football, lacrosse, wrestling, ice hockey and rugby), field and court (basketball, field hockey, soccer, tennis, volleyball, gymnastics, softball and baseball), individual (track and field, swimming, mixed rifle, mixed skiing, golf, dance, diving, cross country, bowling, archery, rowing and cheerleading) and other. Other was used for sport classification when no sport was connected to injury records.

Statistical analyses

With a previously reported high school athlete COVID-19 incidence rate of 5.2 per 1000 athlete-years,²⁹ CR of 2.6 per 100 athlete-years⁸ and the recommended overall model fit of 0.21,³⁰ a total of 418 COVID-19 infections and up to 5 df would be required to reduce risk of model overfitting.³⁰

All data were assessed for missingness prior to analyses (gender, 0%; age, 0%; date of injury, 3%; sport, 1.7%; body part, 1.6%; return to play, <0.1%; and pain, <0.1%), with data demonstrating minimal missingness. Complete case analyses were performed. Participant statistics were described using mean (SD) for continuous, normally distributed variables, median (25th percentile, 75th percentile) for non-normally distributed continuous variables and frequencies (percentages) for categorical variables. Concussion injury incidence rate for athletes who sustained a COVID-19 infection prior to a concussion and those who did not,³¹ with 95% CIs was calculated per 1000 athletes who participated within 1 academic year (1000 athlete-years).³¹ As normality cannot be assumed for incidence rates, point estimates and 95% CIs were logarithmically transformed. All 0 point estimate 95%

Table 1 Participant demographics

	No COVID-19 (n=72 092)	With COVID-19 (n=430)
Age (years)	17.1 (2.7)	17.0 (1.9)
Gender (female), n (%)	32 766 (45)	202 (47)
Sport, n (%)		
Collision	29 558 (41)	133 (31)
Field and court	28 837 (40)	198 (46)
Individual	12 256 (17)	86 (20)
Other	1441 (2)	13 (3)

Results are reported as mean (SD) or count (percentage).

CI were calculated using the exact method.³² Results were stratified by gender, severity and sport. Rate ratios (RR) with 95% CIs were calculated.

Poisson regressions were performed to estimate the rate of concussions following COVID-19 infection. Overdispersion was not present. All models controlled for included important covariables of state, gender, and an offset of the log of COVID-19 infections or athletes who did not sustain a COVID-19 infection in person-years. A cluster at the school level was considered for inclusion in the primary analyses. However, due to the number of athletes who suffered from COVID-19 and then a concussion, it was determined that clustering would only be included as a sensitivity analysis. Potential non-linear associations between factors and outcome were assessed prior to analyses, with all factors demonstrating linear associations. Sensitivity analyses included (1) incorporating high schools as clusters with cluster-robust SEs, (2) stratifying by gender, and (3) comparing athlete concussion risk for those who sustained a bacterial or viral infection that was not a COVID-19 illness and those who did not suffer any illness prior to concussion. All analyses were performed in R V.4.01 (R Foundation for Statistical Computing, Vienna, Austria; <http://www.R-project.org/>) using the *nanjia* package for missingness assessment, *fmsb* package for rate ratio analyses, the *glm* function for Poisson models, the *AER* package to assess for overdispersion and the *GenBinomApps* package for exact method 95% CIs.

RESULTS

General demographics

A total of 72 522 athletes (female, 32 968; male, 39,554) participated in high school sport during the 2020–2021 academic school year (table 1).

General concussion results

High school athletes tested positive for 430 COVID-19 infections during the academic year. A total of 1273 concussions were documented during the school year, resulting in a CR of 17.6 (95% CI 14.8 to 20.8) per 1000 athlete-years. A total of 1241 concussions occurred in athletes who did not previously have a COVID-19 infection and 32 concussions from athletes who did have COVID-19 (table 1). The median time loss from concussion in both groups was 18 (7, 30). The median time from COVID-19 infection to concussion was 14 (7, 22) days. No athletes who tested positive for COVID-19 and then had a concussion sustained a concussion prior to COVID-19 or after 60 days following return from COVID-19. All COVID-19 athletes who suffered a concussion demonstrated COVID-19 symptoms (table 2).

Table 2 Participant concussion rates (number of concussions per 1000 athlete-years) by COVID-19-infected athletes and severity

	Concussion	
	No COVID-19 (n=72 092)	With COVID-19 (n=430)
Overall	17.2 (14.5 to 20.5)	74.4 (55.2 to 100.4)
Minor	4.1 (3.4 to 4.9)	13.9 (10.1 to 19.2)
Moderate	8.9 (7.5 to 10.6)	46.6 (34.2 to 63.3)
Severe	4.3 (3.6 to 5.1)	13.9 (10.2 to 19.2)

All injury rates are reported per 1000 athlete-years with 95% CIs.
Minor injuries are time loss injuries of 1–7 days, moderate=8–27 days, severe=28+ days.

Gender differences in CRs

Girl athletes who sustained a COVID-19 infection were 3.2 (95% CI 1.8 to 5.9) times more likely to suffer from a concussion following return to sport after COVID-19 infection compared with those not sustaining a COVID-19 infection. Boy athletes who sustained a COVID-19 infection were 2.8 (95% CI 1.6 to 4.8) times more likely to suffer from a concussion following return to sport after infection compared with those not sustaining a COVID-19 infection (table 3).

Sport-specific differences in CRs

Collision sport athletes who sustained a COVID-19 infection were 8.9 (95% CI 4.9 to 16.3) times more likely to sustain a concussion following return to sport after infection compared with those collision athletes not sustaining a COVID-19 infection. Field and court sport athletes who sustained a COVID-19 infection were 9.7 (95% CI 5.6 to 17.0) times more likely to suffer from a concussion following return to sport after infection compared with those field and court athletes not sustaining a COVID-19 infection. No individual or other sport athletes suffered a concussion following return to sport after infection (table 4).

Risk of concussion following COVID-19 infection

High school athletes who sustained a COVID-19 infection demonstrated an unadjusted (2.9, 95% CI 1.9 to 4.5, $p<0.001$) and adjusted (3.1, 95% CI 2.0 to 4.7, $p<0.001$) increased rate of concussion following recovery from illness compared with athletes who did not sustain a COVID-19 infection.

Sensitivity analyses

When controlling for clustering at the high school level, athletes who sustained a COVID-19 infection demonstrated (1.6, 95% CI 1.1, 25.9, $p<0.001$) a higher CR following recovery from illness compared with athletes who did not sustain a COVID-19 infection. Female high school athletes who sustained a COVID-19 infection demonstrated (5.7, 95% CI 3.1 to 10.7, $p<0.001$) a higher CR

Table 3 Concussion rates stratified by gender per 1000 athletes-years by COVID-19-infected athletes and severity

	Concussion			
	Girls		Boys	
	No COVID-19 (n=32 766)	With COVID-19 (n=202)	No COVID-19 (n=39 326)	With COVID-19 (n=228)
Overall	15.4 (12.8 to 18.5)	49.5 (34.7 to 70.0)	19.0 (15.9 to 22.8)	52.6 (37.4 to 74.1)
Minor	3.7 (3.0 to 4.5)	14.8 (10.3 to 21.4)	4.4 (3.7 to 5.4)	0 (0 to 10.0)
Moderate	8.0 (6.6 to 9.6)	24.8 (17.3 to 35.5)	9.8 (8.1 to 11.7)	48.2 (34.2 to 60.0)
Severe	3.7 (3.0 to 4.6)	9.9 (0.0 to 23.6)	4.8 (4.0 to 5.8)	4.4 (0.0 to 13.0)

All injury rates are reported per 1000 athlete-years with 95% CIs.
Minor injuries are time loss injuries of 1–7 days, moderate=8–27 days, severe=28+ days.

Table 4 Concussion rates stratified by sport groups per 1000 athlete-years by COVID-19-infected athletes

	Concussion	
	No COVID-19	With COVID-19
Collision	8.5 (7.1 to 10.1)	75.8 (52.3 to 109.8)
Field and court	6.2 (5.2 to 7.4)	60.6 (42.8 to 85.8)
Individual	2.0 (1.7 to 2.5)	0 (0 to 26.7)
Other	0.6 (0.5 to 0.8)	0 (0 to 142.3)

All injury rates are reported per 1000 athlete-years with 95% CIs.

following recovery from illness compared with athletes who did not sustain a COVID-19 infection. Male high school athletes who sustained a COVID-19 infection demonstrated an (3.4, 95% CI 1.9 to 6.1, $p < 0.001$) increased CR following recovery from illness compared with athletes who did not sustain a COVID-19 infection.

Out of the 123 illnesses who were not infected with COVID-19, only one athlete sustained a concussion following recovery from illness. As a result, it was not possible to perform any statistical analyses using these data.

DISCUSSION

The main findings of this study were that in athletes with a COVID-19 infection, when controlling for gender and state, there was a threefold rate of concussion following COVID-19 infection compared with athletes who did not report a COVID-19 infection. Those who identified as girls and boys demonstrated similar CRs following COVID-19. Collision, field and court athletes who reported a COVID-19 infection had ninefold higher rates of concussion following recovery from illness. Only one concussion was sustained following a general infection, suggesting that concussion risk following a general infection may be dissimilar to COVID-19.

High school athletes who reported a previous COVID-19 infection demonstrated greater rates of sustaining a concussion within 60 days following return to sporting activities. The overall count and rates of concussion^{8 16} and COVID-19 infection³³ in these data were similar to previous literature, suggesting the generalisability of these epidemiological findings. Athletes who suffer from an infection can encounter fatigue and chronic respiratory or cardiac symptoms.³⁴ These symptoms can diminish physical activity or sport training,³⁵ potentially increasing injury risk following recovery from illness.²⁹ High school athletes who sustain a COVID-19 infection have reported similar musculoskeletal injury following illness compared with other infections.²⁹ However, a concussion has different mechanisms of injury⁴ and aetiology^{5 6 23} compared with general musculoskeletal injury. Further, within these data, no high school athletes who sustained a concussion following COVID-19 recovery suffered a musculoskeletal injury prior to either COVID-19 infection or concussion. Prior research has observed an association between previous lower extremity injury and increased concussion risk in high school athletes.³⁶ These findings suggest that other mechanisms may impact concussion risk following COVID-19 recovery. Potential explanations include detriments to spatial-temporal cognition,³⁷ dynamic balance deficits³⁸ or increased inflammation.³⁹ However, these are only speculative. Causal investigations are necessary to determine the underlying causes of this risk.

Female and male athletes demonstrated similar CRs following COVID-19 infections. This is in contrast to previous literature where high school female athletes have reported greater CRs compared with comparable sports by both genders, such as

soccer and basketball.^{8 9} Within these data, female athletes did report a higher rate of minor and severe concussions compared with male athletes. High school female athletes are more likely to report concussions compared with male athletes^{12 40} and are far more likely to have severe concussion symptoms.⁴¹

Collision, field and court sports reported similar CRs following COVID-19 infections. While American football has the highest concussion rates for any sport,^{4 16} rates reported for girl athletes are higher than for boy athletes in sports comparable for both genders,^{8 9} with girls' soccer reporting the highest rates.⁸ These discrepancies between genders may influence these sport findings. Individual sports, such as cross country and golf, reported no concussions following COVID-19 infections. While individual sports report lower CRs compared with collision, field and court sport,^{4 16} the absence of concussions following recovery from COVID-19 may be related to the lack of contact with other athletes. However, further research is needed to investigate these potential sport differences.

These results have practical clinical applications. Following recovery from COVID-19 infections, clinicians may consider examining high school athletes for spatial-temporal reasoning,⁴² dynamic balance⁴³ and cardiovascular conditioning.^{44 45} Further, a return to a sport programme involving gradual training and practice loading⁴⁶ to avoid acute spikes in overall sport loading may be considered.⁴⁷ Athletes who participate in sports with high levels of potential head contacts may benefit from limited player-to-player contact initially when returning to sport.⁴⁸

Limitations of this study include the potential for selection bias, as high schools opted into sport participation within the 2020–2021 season. Athlete injury reporting bias may have influenced these results in either direction. There are limited data on individual state COVID-19 exposure prevalence and temporality. Due to the number of concussions sustained by athletes who reported COVID-19, there was insufficient power to control for more covariables. Previous injury data were not collected, which may confound these results. Sport was collapsed for reporting and analysis, which increases the risk of residual confounding. Training and competition minutes or hours of exposure were not collected, decreasing the precision of these findings. States within these data sustained different COVID-19 infection rates, which may influence these findings. There is the potential for sparse data bias, as some data were reported with zero outcomes. Causality cannot be interpreted with these data.

CONCLUSION

Athletes with a COVID-19 infection had higher CRs within 60 days following recovery from infection compared with athletes without an infection. Female and male high school athletes demonstrated similar CRs following COVID-19 infections. Athletes participating in collision, field and court sports were the most vulnerable to concussion following COVID-19 infection. Additional research is needed to understand why athletes with a recent COVID-19 infection are at higher risk of concussion. Clinical judgement or concern may guide further evaluation or a more cautious return to sport in athletes recovering from COVID-19.

Twitter Garrett S Bullock @DRGSBullock and Carolyn A Emery @CarolynAEmery

Contributors GSB, VRN and ES conceived the study idea. GSB, CE, VRN, AP-U and ES were involved in the design and planning. GSB and ES wrote the first draft of the manuscript. GSB, CE, VRN, AP-U, RGG, CT and ES critically revised the manuscript. GSB, CE, VRN, AP-U, RGG, CT and ES approved the final version of the manuscript. ES guarantees the veracity of these findings.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were involved in the design, conduct, reporting or dissemination plans of this research. Refer to the Methods section for further details.

Patient consent for publication Not applicable.

Ethics approval This study obtained favourable ethics approval from the Preferred Reporting Items for Systematic Reviews and Meta-Analyses Health Institutional Review Board (Pro00054926). Informed consent was not required as data used were anonymous.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No data are available. Due to ethical considerations, data is not available.

This article is made freely available for personal use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

ORCID iDs

Garrett S Bullock <http://orcid.org/0000-0003-0236-9015>

Carolyn A Emery <http://orcid.org/0000-0002-9499-6691>

Ellen Shanley <http://orcid.org/0000-0001-8159-8072>

REFERENCES

- Veliz P. Variation in national survey estimates and youth traumatic brain injury—where does the truth lie? *JAMA Pediatr* 2019;173:399:399–99..
- Veliz P, McCabe SE, Eckner JT, et al. Prevalence of concussion among US adolescents and correlated factors. *JAMA* 2017;318:1180–2.
- Castile L, Collins CL, McIlvain NM, et al. The epidemiology of new versus recurrent sports concussions among high school athletes, 2005–2010. *Br J Sports Med* 2012;46:603–10.
- Marar M, McIlvain NM, Fields SK, et al. Epidemiology of concussions among United States high school athletes in 20 sports. *Am J Sports Med* 2012;40:747–55.
- Buzzini SRR, Guskiewicz KM. Sport-Related concussion in the young athlete. *Curr Opin Pediatr* 2006;18:376–82.
- Patel DR, Greydanus DE. Neurologic considerations for adolescent athletes. *Adolesc Med* 2002;13:569–78.
- Denke NJ. Brain injury in sports. *J Emerg Nurs* 2008;34:363–4.
- Schallmo MS, Weiner JA, Hsu WK. Sport and sex-specific reporting trends in the epidemiology of concussions sustained by high school athletes. *J Bone Joint Surg Am* 2017;99:1314–20.
- Gessel LM, Fields SK, Collins CL, et al. Concussions among United States high school and collegiate athletes. *J Athl Train* 2007;42:495–503.
- Baker JG, Leddy JJ, Darling SR, et al. Gender differences in recovery from sports-related concussion in adolescents. *Clin Pediatr (Phila)* 2016;55:771–5.
- Preiss-Farzanegan SJ, Chapman B, Wong TM, et al. The relationship between gender and postconcussion symptoms after sport-related mild traumatic brain injury. *PM&R* 2009;1:245–53.
- Miyashita TL, Diakogeorgiou E, VanderVegt C. Gender differences in concussion reporting among high school athletes. *Sports Health* 2016;8:359–63.
- Covassin T, Elbin RJ. The female athlete: the role of gender in the assessment and management of sport-related concussion. *Clin Sports Med* 2011;30:125–31.
- Ono KE, Burns TG, Bearden DJ, et al. Sex-based differences as a predictor of recovery trajectories in young athletes after a sports-related concussion. *Am J Sports Med* 2016;44:748–52.
- Harmon KG, Drezner J, Gammons M, et al. American medical Society for sports medicine position statement. *Clinical Journal of Sport Medicine* 2013;23:1–18.
- Marshall SW, Guskiewicz KM, Shankar V, et al. Epidemiology of sports-related concussion in seven us high school and collegiate sports. *Inj Epidemiol* 2015;2:13.
- Maltezou HC, Magaziotou I, Dedoukou X, et al. Children and adolescents with SARS-cov-2 infection: epidemiology, clinical course and viral loads. *Pediatr Infect Dis J* 2020;39:e388–92.
- Zachariah P, Johnson CL, Halabi KC, et al. Epidemiology, clinical features, and disease severity in patients with coronavirus disease 2019 (COVID-19) in a children's hospital in new york city, new york. *JAMA Pediatr* 2020;174.
- Crook H, Raza S, Nowell J, et al. n.d. Long covid—mechanisms, risk factors, and management. *BMJ*:1648.
- De Sousa RAL, Impropita-Caria AC, Aras-Júnior R, et al. Physical exercise effects on the brain during COVID-19 pandemic: links between mental and cardiovascular health. *Neurol Sci* 2021;42:1325–34.
- Raony I, de Figueiredo CS, Pandolfo P, et al. n.d. Psycho-neuroendocrine-immune interactions in COVID-19: potential impacts on mental health. *Front Immunol*;11:1170.
- Zimmermann P, Pittet LF, Curtis N. How common is long COVID in children and adolescents? *Pediatric Infectious Disease Journal* 2021;40:e482–7.
- Guskiewicz KM, Bruce SL, Cantu RC, et al. National athletic trainers' association position statement: management of sport-related concussion. *J Athl Train* 2004;39:280–97.
- Bahr R, Clarsen B, Derman W, et al. International olympic committee consensus statement: methods for recording and reporting of epidemiological data on injury and illness in sports 2020 (including the STROBE extension for sports injury and illness surveillance (STROBE-SIIS)). *Orthop J Sports Med* 2020;8:2325967120902908.
- Ljungqvist A, Jenoure P, Engebretsen L, et al. The International Olympic Committee (IOC) consensus statement on periodic health evaluation of elite athletes March 2009. *British Journal of Sports Medicine* 2009;43:631–43.
- Rae K, Orchard J. The orchard sports injury classification system (OSICS) version 10. *Clin J Sport Med* 2007;17:201–4.
- McCrory P, Meeuwisse W, Dvořák J, et al. Consensus statement on concussion in sport—the 5th international conference on concussion in sport held in berlin, october 2016. *Br J Sports Med* 2017;51:838–47.
- Powell JW, Barber-Foss KD. Sex-Related injury patterns among selected high school sports. *Am J Sports Med* 2000;28:385–91.
- Shanley E, Thigpen CA, Prats-Urbe A, et al. Subsequent musculoskeletal injury incidence are similar between general infection and COVID-19 among high school athletes in the united states despite higher illness incidence during academic years 2019–2020 and 2020–2021. *J Orthop Sports Phys Ther* 2022;52:546–53.
- Riley RD, Debray TPA, Collins GS, et al. Minimum sample size for external validation of a clinical prediction model with a binary outcome. *Stat Med* 2021;40:4230–51.
- Knowles SB, Marshall SW, Guskiewicz KM. Issues in estimating risks and rates in sports injury research. *J Athl Train* 2006;41:207–15.
- Vollset SE. Confidence intervals for a binomial proportion. *Stat Med* 1993;12:809–24.
- Leidman E, Duca LM, Omura JD, et al. COVID-19 trends among persons aged 0–24 years—United States, March 1–december 12, 2020. *MMWR Morb Mortal Wkly Rep* 2021;70:88–94.
- Carfi A, Bernabei R, Landi F, et al. Persistent symptoms in patients after acute COVID-19. *JAMA* 2020;324:603–5.
- Constandt B, Thibaut E, De Bosscher V, et al. Exercising in times of lockdown: an analysis of the impact of COVID-19 on levels and patterns of exercise among adults in Belgium. *Int J Environ Res Public Health* 2020;17:4144:11..
- Lutz A, Thigpen C, Gilliland RG, et al. Lower extremity musculoskeletal injury is associated with increased risk of concussion in high school athletes. *Orthopaedic Journal of Sports Medicine* 2020;8(7_suppl6):232596712050039.
- Howell DR, Osternig LR, Chou L-S. Single-task and dual-task tandem gait test performance after concussion. *J Sci Med Sport* 2017;20:622–6.
- Martinez-Córcoles V, Nieto-Gil P, Ramos-Petersen L, et al. Balance performance analysis after the COVID-19 quarantine in children aged between 8 and 12 years old: longitudinal study. *Gait Posture* 2022;94:203–9.
- Cothran TP, Kellman S, Singh S, et al. A brewing storm: the neuropsychological sequelae of hyperinflammation due to COVID-19. *Brain Behav Immun* 2020;88:957–8.
- McCrea M, Hammeke T, Olsen G, et al. Unreported concussion in high school football players: implications for prevention. *Clin J Sport Med* 2004;14:13–7.
- Tanveer S, Zecavati N, Delasobera EB, et al. Gender differences in concussion and postinjury cognitive findings in an older and younger pediatric population. *Pediatric Neurology* 2017;70:44–9.
- Howell DR, Lynall RC, Buckley TA, et al. Neuromuscular control deficits and the risk of subsequent injury after a concussion: a scoping review. *Sports Med* 2018;48:1097–115.
- Davis GA, Anderson V, Babl FE, et al. What is the difference in concussion management in children as compared with adults? A systematic review. *Br J Sports Med* 2017;51:949–57.
- Schneider KJ, Leddy JJ, Guskiewicz KM, et al. Rest and treatment/rehabilitation following sport-related concussion: a systematic review. *Br J Sports Med* 2017;51:930–4.
- Powell C, McCaulley B, Scott Brosky Z, et al. The effect of aerobic exercise on adolescent athletes post-concussion: a systematic review and meta-analysis. *Int J Sports Phys Ther* 2020;15:650–8.
- Miutz LN, Burma JS, Lapointe AP, et al. Physical activity following sport-related concussion in adolescents: a systematic review. *J Appl Physiol (1985)* 2022;132:1250–66.
- Purcell LK, Davis GA, Gioia GA. What factors must be considered in “return to school” following concussion and what strategies or accommodations should be followed? A systematic review. *Br J Sports Med* 2019;53:250–50.
- Emery CA, Black AM, Kolstad A, et al. What strategies can be used to effectively reduce the risk of concussion in sport? A systematic review. *Br J Sports Med* 2017;51:978–84.